arXiv:1407.1284v2 [math.SP] 16 Jul 2014

ON THE ESSENTIAL SPECTRUM OF N-BODY SYSTEMS WITH ASYMPTOTICALLY HOMOGENEOUS OF ORDER ZERO INTERACTIONS

VLADIMIR GEORGESCU AND VICTOR NISTOR

ABSTRACT. We overview some of our recent results on the essential spectrum of N-body Hamiltonians with potentials defined by functions that have radial limits at infinity. The results extend the HVZ theorem which describes the essential spectrum of usual N-body Hamiltonians. The proof is based on a careful study of algebras generated by potentials and their cross-products. We also describe the topology on the spectrum of these algebras, thus extending to our setting a result of A. Mageira. Our techniques apply to more general classes of potentials associated to translation invariant algebras of bounded uniformly continuous functions on a finite dimensional vector space X.

INTRODUCTION

Let X be a finite dimensional real vector space and, for each linear subspace Y of X, let $V_Y : X/Y \to \mathbb{R}$ be a Borel function. To simplify the statements, we assume $V_Y = 0$, except for a finite number of Y. We keep the notation V_Y for the function on X given by $V_Y \circ \pi_Y$, where $\pi_Y : X \to X/Y$ is the natural map. In this paper, we use crossed-products of C^*-algebras to study the essential spectrum of Hamiltonians of the form

$$H := h(P) + \textstyle\sum_Y V_Y \,, \tag{1}$$

under certain conditions on the potentials V_Y. Here $h : X^* \to [0, \infty[$ is a continuous, proper function and P is the momentum observable (recall that *proper* means that $\lim_{|k|\to\infty} h(k) = +\infty$). More precisely, $h(P) = \mathcal{F}^{-1} M_h \mathcal{F}$, where $\mathcal{F} : L^2(X) \to L^2(X^*)$ is the Fourier transform and M_h is the operator of multiplication by h (formally $P = -\mathrm{i}\nabla$).

Operators of this form cover the Hamiltonians that are currently the most interesting (from a physical point of view) Hamiltonians of N-body systems. Here are two main examples. In a generalized version of the non-relativistic case, a scalar product is given on X, so, by taking $h(\xi) = |\xi|^2$, we get $h(P) = \Delta$, the positive Laplacian. In the simplest relativistic case, $X = (\mathbb{R}^3)^N$ and, writing the momentum $P = (P_1, \ldots, P_N)$, we have $h(P) = \sum_{k=1}^N (P_k^2 + m_k^2)^{1/2}$ for some real numbers m_k. We refer to [3] for a thorough introduction to the subject and study of these systems.

To state our results, we need some notations and a definition. Let $\mathbb{S}_X := (X \smallsetminus \{0\})/\mathbb{R}_+$ be *the sphere at infinity of* X, i.e. the set of all half-lines $\widehat{a} := \mathbb{R}_+ a$ with $\mathbb{R}_+ =]0, \infty[$ and $a \in X \setminus \{0\}$. A function $v : X \to \mathbb{C}$ is said *to have uniform radial limits at infinity* if $v(\widehat{a}) := \lim_{r\to\infty} v(ra)$ exists uniformly in $\widehat{a} \in \mathbb{S}_X$. Clearly then $v(\widehat{a}) = \lim_{r\to\infty} v(ra + x)\,,\ \forall x \in X$. More generally, we are interested in functions v such that $\lim_{r\to\infty} v(ra + x)$ exists for all $x \in X$. The limit may depend on x and defines a function $\tau_\alpha(v) : X \to \mathbb{C}$, where $\alpha := \widehat{a}$. For example, let us consider $v = V_Y$. Then $\tau_\alpha(V_Y)(x) = \lim_{r\to\infty} V_Y(r\pi_Y(a) + \pi_Y(x))$. In particular, $\tau_\alpha(V_Y) = V_Y$ whenever $\alpha := \widehat{a} \subset Y$ (i. e. $a \in Y$). On the other hand, if $V_Y : X/Y \to \mathbb{C}$ has uniform radial limits at infinity and $\widehat{a} = \alpha \not\subset Y$, then $\pi_Y(\alpha) := \mathbb{R}_+ \pi_Y(a) = \widehat{a + Y} \in \mathbb{S}_{X/Y}$ is well defined and $\tau_\alpha(V_Y)(x) = V_Y(\pi_Y(\alpha))$ turns out to be a *constant*. We denote by $\overline{\cup}_\alpha A_\alpha := \overline{\cup_\alpha A_\alpha}$ the closure of the union of a given family of sets. Here is the main result of the paper:

Theorem 1. *Assume that each of the functions $V_Y : X/Y \to \mathbb{R}$ is bounded and has uniform radial limits at infinity. For each $\alpha \in \mathbb{S}_X$, we denote*

$$\tau_\alpha(H) = h(P) + \sum_Y \tau_\alpha(V_Y) = h(P) + \sum_{Y \supset \alpha} V_Y + \sum_{Y \not\supset \alpha} V_Y(\pi_Y(\alpha))\,. \tag{2}$$

Then $\sigma(\tau_\alpha(H)) = [c_\alpha, \infty)$ for some real c_α and $\sigma_{\mathrm{ess}}(H) = \overline{\cup}_{\alpha\in\mathbb{S}_X} \sigma(\tau_\alpha(H)) = [\inf_\alpha c_\alpha, \infty)$.

Unbounded potentials are considered in Theorem 4. If all the radial limits are zero, which is the case of the usual N-body potentials, then the terms corresponding to $\alpha \not\subset Y$ are dropped in (2). Consequently, if $h(P) = \Delta$ is the non-relativistic kinetic energy, we recover the Hunziker, van Winter, Zhislin (HVZ) theorem. Descriptions of the essential spectrum of various classes of self-adjoint operators in terms of limits at infinity of translates of the operators

have already been obtained before, see for example [8, 12] and [9] (in historical order). Our approach is based on the "localization at infinity" technique developed in [6, 7] in the context of crossed-product C^*-algebras.

Let us sketch the main idea of this approach. Let $\mathcal{A}$ be a translation invariant C^*-algebra of bounded uniformly continuous functions containing the functions which have a limit at infinity and let $\widehat{\mathcal{A}}$ be its character space. Note that $\widehat{\mathcal{A}}$ is a compact topological space that naturally contains X as an open dense subset and $\delta(\mathcal{A}) = \widehat{\mathcal{A}} \setminus X$ can be thought as a boundary of X at infinity. Then to each self-adjoint operator H affiliated[1] to the crossed product $\mathcal{A} \rtimes X$ of $\mathcal{A}$ by the action of X one may associate a family of self-adjoint operators $H_\varkappa$ affiliated to $\mathcal{A} \rtimes X$ indexed by the characters $\varkappa \in \delta(\mathcal{A})$. This family completely describes the image of H (in the sense of affiliated operators) in the quotient of $\mathcal{A} \rtimes X$ with respect to the ideal of compact operators. In particular, the essential spectrum of H is the closure of the union of the spectra of the operators $H_\varkappa$. These operators are the *localizations at infinity of* H, more precisely, $H_\varkappa$ *is the localization of* H *at the point* $\varkappa$.

Once chosen the algebra $\mathcal{A}$, in order to use these techniques of this paper concrete situations, one still needs the following: (1) to have a good description of the character space of the abelian algebra $\mathcal{A}$, and (2) to have an efficient criterion for affiliation to the crossed product $\mathcal{A} \rtimes X$. In this paper, we explain also how to achieve these two steps.

1. Crossed products and localization at infinity

We begin with some general facts. Let $\mathcal{C}_{\mathrm{b}}(X)$ be the algebra of bounded continuous functions, $\mathcal{C}_0(X)$ the ideal of functions vanishing at infinity, $\mathcal{C}(X^+) = \mathbb{C} + \mathcal{C}_0(X)$, and $\mathcal{C}^{\mathrm{u}}_{\mathrm{b}}(X)$ the subalgebra of bounded uniformly continuous functions on X. If $p \in X^*$ and $q \in X$ then we define unitary operators S_p and T_q on $L^2(X)$ by

$$(S_p f)(x) \,=\, \mathrm{e}^{\mathrm{i}p(x)} f(x) \quad \text{ and } \quad (T_q f)(q) \,=\, f(x+q)\,. \tag{3}$$

One sometimes writes $p(x) = \langle x|p\rangle$. The symbol $A^{(*)}$ used below means that the relation should hold for A and A^*.

Definition 1. *We say that a bounded operator A on $L^2(X)$* has the position-momentum limit property *if*

$$\lim_{p\to 0} \|[S_p, A]\| \,=\, 0 \quad \text{ and } \quad \lim_{q\to 0} \|(T_q - 1)A^{(*)}\| \,=\, 0\,.$$

It is clear that the set of operators which have the position-momentum limit property is a C^*-algebra. This algebra is equal to the (image in $\mathcal{B}(L^2(X))$ of the crossed products $\mathcal{C}^{\mathrm{u}}_{\mathrm{b}}(X) \rtimes X$ (see [6]). More precisely, if $\mathcal{A}$ is an arbitrary translation invariant C^*-subalgebra of $\mathcal{C}^{\mathrm{u}}_{\mathrm{b}}(X)$, then there is a natural realization of the abstract crossed product $\mathcal{A} \rtimes X$ as a C^*-algebra of operators on $L^2(X)$ and we do not distinguish the two algebras. To describe this concrete version of $\mathcal{A} \rtimes X$, and for later use, we need some more notations.

If $\varphi : X \to \mathbb{C}$ and $\psi : X^* \to \mathbb{C}$ are measurable functions then $\varphi(Q)$ and $\psi(P)$ are the operators on $L^2(X)$ defined as follows: $\varphi(Q) := M_\phi$ acts as multiplication by φ and $\psi(P) = \mathcal{F}^{-1} M_\psi \mathcal{F}$, where $\mathcal{F}$ is the Fourier transform $L^2(X) \to L^2(X^*)$ and M_ψ is the operator of multiplication by ψ. The group C^*-algebra of X, i.e. the closed subspace of $B(L^2(X))$ generated by convolution with continuous functions with compact support, will be denoted $C^*(X)$. The map $\psi \mapsto \psi(P)$ is an isomorphism between $\mathcal{C}_0(X^*)$ and $C^*(X)$. With these notations, we have: $\mathcal{A} \rtimes X$ is the norm closed linear space of bounded operators on $L^2(X)$ generated by the products $\varphi(Q)\psi(P)$ with $\varphi \in \mathcal{A}$ and $\psi \in \mathcal{C}_0(X^*)$. In particular, $\mathcal{A} \rtimes X$ consists of operators that have the position-momentum limit property.

Our next goal is to recall the definition of localizations at infinity for operators that have the position-momentum limit property. Let us fix then an algebra $\mathcal{A}$ such that $\mathcal{C}(X^+) \subset \mathcal{A}$. As we explained in the introduction, $\widehat{\mathcal{A}}$ is a compactification of X and $\delta(\mathcal{A}) = \widehat{\mathcal{A}} \setminus X$ is a compact set which can be characterized as the set of characters $\varkappa$ such that $\varkappa|_{\mathcal{C}_0(X)} = 0$. The largest $\mathcal{A}$ that we may consider is $\mathcal{C}^{\mathrm{u}}_{\mathrm{b}}(X)$ and we denote $\gamma(X) := \widehat{\mathcal{C}^{\mathrm{u}}_{\mathrm{b}}(X)}$ its character space and $\delta(X) = \gamma(X) \setminus X$. For a general $\mathcal{A}$ now, the map $\varkappa \mapsto \varkappa|_{\mathcal{A}}$ is a continuous surjective map $\delta(X) \to \delta(\mathcal{A})$, and hence $\delta(\mathcal{A})$ is a quotient of the compact space $\delta(X)$. If $q \in X$ and φ is a function on X then $T_q\varphi$ is its translation by q, defined as in (3). We extend now this definition of T_q by replacing $q \in X$ with a character $\varkappa \in \widehat{\mathcal{A}}$:

$$(T_\varkappa \varphi)(x) = \varkappa(T_x \varphi)\,, \quad \text{ for any } \ \varphi \in \mathcal{A}\,, \ \ \varkappa \in \widehat{\mathcal{A}} \ \text{ and, } \ x \in X. \tag{4}$$

[1] A self-adjoint operator H on a Hilbert space $\mathcal{H}$ is said to be *affiliated* to a C^*-algebra $\mathscr{A}$ of operators on $\mathcal{H}$ if one has $(H + \mathrm{i})^{-1} \in \mathscr{A}$. Then $\varphi(H) \in \mathscr{A}$ for any $\varphi \in \mathcal{C}_0(\mathbb{R})$.

It is clear that $T_\varkappa\varphi \in \mathcal{C}_{\rm b}^{\rm u}(X)$ and that its definition coincides with the previous one if $\varkappa = q \in X$. Moreover, we also get "translations at infinity" of $\varphi \in \mathcal{A}$ by elements $\varkappa \in \delta(\mathcal{A})$; note however that such a translation does not belong to $\mathcal{A}$ in general. Also, the function $\varkappa \mapsto T_\varkappa\varphi \in \mathcal{C}_{\rm b}^{\rm u}(X)$ defined on $\widehat{\mathcal{A}}$ is continuous if $\mathcal{C}_{\rm b}^{\rm u}(X)$ is equipped with the topology of local uniform convergence, hence $T_\varkappa\varphi = \lim_{q\to\varkappa} T_q\varphi$ in this topology for any $\varkappa \in \delta(\mathcal{A})$. If A is an operator on $L^2(X)$, let $\tau_q(A) = T_qAT_q^*$ be its translation by $q \in X$. Clearly $\tau_q(\varphi(Q)) = (T_q\varphi)(Q)$. If $A \in \mathcal{A} \rtimes X$, then we may also consider "translations at infinity" by elements of the boundary $\delta(\mathcal{A})$ of X in $\widehat{\mathcal{A}}$ and we get a useful characterization of the compact operators. The following facts are mainly consequences of [7, Theorem 1.15]:

- If $\varkappa \in \widehat{\mathcal{A}}$, then there is a unique morphism $\tau_\varkappa : \mathcal{A} \rtimes X \to \mathcal{C}_{\rm b}^{\rm u}(X) \rtimes X$ such that
$$\tau_\varkappa(\varphi(Q)\psi(P)) \;=\; (T_\varkappa\varphi)(Q)\psi(P)\,, \quad \text{for all } \varphi \in \mathcal{C}_{\rm b}^{\rm u}(X),\, \psi \in \mathcal{C}_0(X)\,.$$
- If $A \in \mathcal{A} \rtimes X$ then $\varkappa \mapsto \tau_\varkappa(A)$ is a strongly continuous map $\widehat{\mathcal{A}} \to \mathcal{B}(L^2(X))$.
- $\cap_{\varkappa\in\delta(\mathcal{A})} \ker\tau_\varkappa = \mathcal{C}_0(X) \rtimes X \equiv \mathscr{K}(X) =$ ideal of compact operators on $L^2(X)$.
- The map $\tau(A) = (\tau_\varkappa(A))_{\varkappa\in\delta(\mathcal{A})}$ induces an injective morphism
$$\mathcal{A} \rtimes X/\mathscr{K}(X) \hookrightarrow \textstyle\prod_{\varkappa\in\delta(\mathcal{A})} \mathscr{C}(X). \tag{5}$$
- If $A \in \mathcal{A} \rtimes X$ is a normal operator then $\sigma_{\rm ess}(A) = \overline{\cup}_{\varkappa\in\delta(\mathcal{A})}\sigma(\tau_\varkappa(A))$.
- If H is a self-adjoint operator on $L^2(X)$ affiliated to $\mathcal{A}$ then for each $\varkappa \in \delta(\mathcal{A})$ the limit $\tau_\varkappa(H) := \text{s-}\lim_{q\to\varkappa} T_qHT_q^*$ exists and $\sigma_{\rm ess}(H) = \overline{\cup}_{\varkappa\in\delta(\mathcal{A})}\sigma(\tau_\varkappa(H))$.

To be precise, the last strong limit means: $\tau_\varkappa(H)$ is a self-adjoint operator (not necessarily densely defined) on $L^2(X)$ and $\text{s-}\lim_{q\to\varkappa}\theta(T_qHT_q^*) = \theta(\tau_\varkappa(H))$ for all $\theta \in \mathcal{C}_0(\mathbb{R})$. It is clear that in the last three statements above one may replace $\delta(\mathcal{A})$ by a subset π if for each $A \in \mathcal{A} \rtimes X$ we have: $\tau_\varkappa(A) = 0\,\forall\varkappa \in \pi \Rightarrow \tau_\varkappa(A) = 0\,\forall\varkappa \in \delta(\mathcal{A})$. In the case of groupoid (pseudo)differential algebras (that is, when $\widehat{\mathcal{A}}$ is a manifold with corners), the morphisms $\tau_\varkappa$ can be defined using restrictions to fibers, as in [10], and the last three statements above remain valid.

2. The Two-Body Case

As a warm up and in order to introduce some general notation, we treat first the two-body case, where complete results may be obtained by direct arguments. The algebra of interactions in the standard two-body case is $\mathcal{C}(X^+)$, and hence the Hamiltonian algebra is

$$\mathcal{C}(X^+) \rtimes X = \mathbb{C} \rtimes X + \mathcal{C}_0(X) \rtimes X = C^*(X) + \mathscr{K}(X) \tag{6}$$

where the sums are direct. Thus $\mathcal{C}(X^+) \rtimes X/\mathscr{K}(X) = C^*(X)$ which finishes the theory. Another elementary case, which has been considered as an example in [6], is $X = \mathbb{R}$ with $\mathcal{C}(\mathbb{R}^+)$ replaced by the algebra $\mathcal{C}(\overline{\mathbb{R}})$ of continuous functions which have limits (distinct in general) at $\pm\infty$. Then there is no natural direct sum decomposition of $\mathcal{C}(\overline{\mathbb{R}}) \rtimes \mathbb{R}$ as in (6), but one has by standard arguments

$$\mathcal{C}(\overline{\mathbb{R}}) \rtimes \mathbb{R}/\mathscr{K}(\mathbb{R}) \simeq C^*(\mathbb{R}) \oplus C^*(\mathbb{R}). \tag{7}$$

Our purpose in this section is to extend (7) to arbitrary X and the main result is an explicit description of the crossed product algebra $\mathcal{C}(\overline{X}) \rtimes X$.

Let $\mathcal{C}(\overline{X})$ be the closure in $\mathcal{C}_{\rm b}(X)$ of the subalgebra of functions homogeneous of degree zero outside a compact set. Then $\mathcal{C}(\overline{X}) = \{u \in \mathcal{C}(X) \mid \lim_{\lambda\to+\infty} u(\lambda a) \text{ exists uniformly in } \widehat{a} \in \mathbb{S}_X\}$, where, we recall, $\widehat{a} := \mathbb{R}_+a$ and $\mathbb{S}_X := (X \smallsetminus \{0\})/\mathbb{R}_+$, so $\widehat{a} \in \mathbb{S}_X$. As a set, the character space of $\mathcal{C}(\overline{X})$ can be identified with the disjoint union $\overline{X} = X \cup \mathbb{S}_X$. The topology induced by the character space on X is the usual one and the intersections with X of the neighbourhoods of some $\alpha \in \mathbb{S}_X$ are the sets which contain a truncated cone C such that there is $a \in \alpha$ such $\lambda a \in C$ if $\lambda \geq 1$. The set of such subsets is a filter $\widetilde{\alpha}$ on X and, if Y is a Hausdorff space and $u : X \to Y$, then $\lim_{\widetilde{\alpha}} u = y$ means that $u^{-1}(V) \in \widetilde{\alpha}$ for any neighborhood V of u. We shall write $\lim_{x\to\alpha} u(x)$ instead of $\lim_{\widetilde{\alpha}} u$. We have that $\mathcal{C}(\overline{X})$ is a translation invariant C^*-subalgebra of $\mathcal{C}_{\rm b}^{\rm u}(X)$ and so the crossed product $\mathcal{C}(\overline{X}) \rtimes X$ is well defined. We have the following explicit description of this algebra.

Proposition 1. *The algebra $\mathcal{C}(\overline{X}) \rtimes X$ acting on $L^2(X)$ consists of bounded operators A that have the position-momentum limit property and are such that the limit $\tau_\alpha(A) = \text{s-lim}_{a\to\alpha} T_a A T_a^*$ exists for each $\alpha = \widehat{a} \in \mathbb{S}_X$. If $A \in \mathcal{C}(\overline{X}) \rtimes X$ and $\alpha \in \mathbb{S}_X$, then $\tau_\alpha(A) \in C^*(X)$ and $\tau(A) : \alpha \mapsto \tau_\alpha(A)$ is norm continuous. The map $\tau : \mathcal{C}(\overline{X}) \rtimes X \to C(\mathbb{S}_X) \otimes C^*(X)$ is a surjective morphism whose kernel is the set of compact operators on $L^2(X)$, which gives*

$$\mathcal{C}(\overline{X}) \rtimes X/\mathscr{K}(X) \cong C(\mathbb{S}_X) \otimes C^*(X). \tag{8}$$

If H is a self-adjoint operator affiliated to $\mathcal{C}(\overline{X}) \rtimes X$ then $\tau_\alpha(H) = \text{s-lim}_{a\to\alpha} T_a H T_a^*$ exists for all $\alpha \in \mathbb{S}_X$ and $\sigma_{\text{ess}}(H) = \cup_\alpha \sigma(\tau_\alpha(H))$.

Equations (7) and (8) follow also from standard properties of crossed-product algebras by amenable groups.

In the next two example $H = h(P) + V$ with $h : X^* \to [0, \infty[$ continuous and proper. We denote by $|\cdot|$ a fixed norm on X^* and by $\mathcal{H}^s$ we denote the usual Sobolev spaces on X ($s \in \mathbb{R}$).

Example 1. Let V be a bounded symmetric operator satisfying: (1) $\lim_{p\to 0} \|[S_p, V]\| = 0$ and (2) the limit $\tau_\alpha(V) = \text{s-lim}_{a\to\alpha} T_a V T_a^*$ exists for each $\alpha \in \mathbb{S}_X$. Then H is affiliated to $\mathcal{C}(\overline{X}) \rtimes X$ and $\tau_\alpha(H) = h(P) + \tau_\alpha(V)$. Moreover, if V is a function then $\tau_\alpha(V)$ is a number, but in general we have $\tau_\alpha(V) = v_\alpha(P)$ for some function $v_\alpha \in \mathcal{C}_{\text{b}}^{\text{u}}(X^*)$.

Example 2. Assume that h is locally Lipschitz and that there exist $c, s > 0$ such that, for all p with $|p| > 1$,

$$|\nabla h(p)| \le c\big(1 + h(p)\big) \quad \text{and} \quad c^{-1}|p|^s \le \big(1 + h(p)\big)^{1/2} \le c|p|^s. \tag{9}$$

Let $V : \mathcal{H}^s \to \mathcal{H}^{-s}$ such that $\pm V \le \mu h(P) + \nu$ for some numbers μ, ν with $\mu < 1$ and satisfying the next two conditions: (1) $\lim_{p\to 0} \|[S_p, V]\|_{\mathcal{H}^s \to \mathcal{H}^{-s}} = 0$, (2) $\forall \alpha \in \mathbb{S}_X$ the limit $\tau_\alpha(V) = \text{s-lim}_{a\to\alpha} T_a V T_a^*$ exists strongly in $\mathcal{B}(\mathcal{H}^s, \mathcal{H}^{-s})$. Then $h(P) + V$ and $h(P) + \tau_\alpha(V)$ are symmetric operators $\mathcal{H}^s \to \mathcal{H}^{-s}$ which induce self-adjoint operators H and $\tau_\alpha(H)$ in $L^2(X)$ affiliated to $\mathcal{C}(\overline{X}) \rtimes X$ and $\sigma_{\text{ess}}(H) = \cup_\alpha \sigma(\tau_\alpha(H))$.

3. N-BODY CASE

We first indicate a general way of constructing N-body Hamiltonians. For each linear subspace $Y \subset X$ let $\mathcal{A}(X/Y) \subset \mathcal{C}_{\text{b}}^{\text{u}}(X/Y)$ be a translation invariant C^*-subalgebra containing $\mathcal{C}_0(X/Y)$ with $\mathcal{A}(X/X) = \mathcal{A}(0) = \mathbb{C}$. We embed $\mathcal{A}(X/Y) \subset \mathcal{C}_{\text{b}}^{\text{u}}(X)$ as usual by identifying v with $v \circ \pi_Y$. Then the C^*-algebra $\mathcal{A}$ generated by these algebras is a translation invariant C^*-subalgebra of $\mathcal{C}_{\text{b}}^{\text{u}}(X)$ containing $\mathcal{C}(X^+)$ and thus we may consider the crossed product $\mathcal{A} \rtimes X$ which is equal to the C^*-algebra generated by the crossed products $\mathcal{A}(X/Y) \rtimes X$. The operators affiliated to $\mathcal{A} \rtimes X$ are N-body Hamiltonians.

The standard N-body algebra corresponds to the minimal choice $\mathcal{A}(X/Y) = \mathcal{C}_0(X/Y)$ and has remarkable properties which makes its study relatively easy (it is graded by the lattice of subspaces of X). Our purpose in this paper is to study what could arguably be considered to be the simplest extension of the classical N-body setting: namely, it is obtained by choosing $\mathcal{A}(X/Y) = \mathcal{C}(\overline{X/Y})$ for all Y. The next more general case would correspond to the choice $\mathcal{A}(X/Y) = \mathcal{V}(X/Y)$ which is the algebra of functions with vanishing oscillating at infinity (the closure in $\sup$ norm of the set of bounded functions of class C^1 with derivatives tending to zero at infinity).

Definition 2. *Let $\mathcal{E}(X)$ be the C^*-subalgebra of $\mathcal{C}_{\text{b}}^{\text{u}}(X)$ generated by $\cup_Y \mathcal{C}(\overline{X/Y})$.*

Clearly $\mathcal{E}(X)$ is a translation invariant C^*-subalgebra of $\mathcal{C}_{\text{b}}^{\text{u}}(X)$ containing $\mathcal{C}(X^+) := \mathcal{C}_0(X) + \mathbb{C}$. If Y is a linear subspace of X then the C^*-algebra $\mathcal{E}(X/Y) \subset \mathcal{C}_{\text{b}}^{\text{u}}(X/Y)$ is well defined and naturally embedded in $\mathcal{E}(X)$: it is the C^*-algebra generated by $\cup_{Z\supset Y} C(\overline{X/Z})$. We have

$$\mathbb{C} = \mathcal{E}(0) = \mathcal{E}(X/X) \subset \mathcal{E}(X/Y) \subset \mathcal{E}(X/Z) \subset \mathcal{E}(X). \tag{10}$$

If $\alpha \in \mathbb{S}_X$, we shall denote by abuse of notation X/α be the quotient $X/[\alpha]$ of X by the subspace $[\alpha] := \mathbb{R}\alpha$ generated by α and let us set $\pi_\alpha = \pi_{[\alpha]}$. It is clear that $\tau_\alpha(u)(x) = \lim_{r\to+\infty} u(ra + x)$ exists $\forall u \in \mathcal{E}(X)$ and that the resulting function $\tau_\alpha(u)$ belongs to $\mathcal{E}(X)$. The map τ_α is an endomorphism of $\mathcal{E}(X)$ and a linear projection of $\mathcal{E}(X)$ onto the C^*-subalgebra $\mathcal{E}(X/\alpha)$.

If $\alpha \in \mathbb{S}_X$ and $\beta \in \mathbb{S}_{X/\alpha}$, then β generates a one dimensional linear subspace $[\beta] := \mathbb{R}\beta \subset X/\alpha$, as above, and hence $\pi_\alpha^{-1}([\beta])$ is a two dimensional subspace of X that we shall denote $[\alpha, \beta]$. We shall identify $(X/\alpha)/\beta$ with

$X/[\alpha,\beta]$. Then we have two idempotent morphisms $\tau_\alpha : \mathcal{E}(X) \to \mathcal{E}(X/\alpha)$ and $\tau_\beta : \mathcal{E}(X/\alpha) \to \mathcal{E}(X/[\alpha,\beta])$. Thus $\tau_\beta\tau_\alpha : \mathcal{E}(X) \to \mathcal{E}(X/[\alpha,\beta])$ is an idempotent morphism. This construction extends in an obvious way to families $\overrightarrow{\alpha} = (\alpha_1,\dots,\alpha_n)$ with $n \leq \dim X$ and $\alpha_1 \in \mathbb{S}_X, \alpha_2 \in \mathbb{S}_{X/\alpha_1}, \alpha_3 \in \mathbb{S}_{X/[\alpha_1,\alpha_2]}, \dots$(we allow $n = 0$ and denote A the set of all such families). The endomorphism $\tau_{\overrightarrow{\alpha}}$ of $\mathcal{E}(X)$ is defined by induction: $\tau_{\overrightarrow{\alpha}} = \tau_{\alpha_n}\dots\tau_{\alpha_1}$. We also define $[\alpha_1,\alpha_2,\dots,\alpha_n]$ by induction, so this is an n-dimensional subspace of X associated to $(\alpha_1,\alpha_2,\dots,\alpha_n)$ and we denote $X/\overrightarrow{\alpha}$ the quotient of X with respect to it. Thus $\tau_{\overrightarrow{\alpha}}$ is an endomorphism of $\mathcal{E}(X)$ and a projection of $\mathcal{E}(X)$ onto the subalgebra $\mathcal{E}(X/\overrightarrow{\alpha})$. The next description of the spectrum of $\mathcal{E}(X)$ extends some results of A. Mageira [11].

Theorem 2. *If $\overrightarrow{\alpha} \in A$ and $a \in X/\overrightarrow{\alpha}$, then $\varkappa(u) = (\tau_{\overrightarrow{\alpha}}u)(a)$ defines a character of $\mathcal{E}(X)$. Conversely, each character of $\mathcal{E}(X)$ is of this form.*

Remark 1. A natural abelian C^*-algebra in the present context is the set $\mathcal{R}(X)$ of all bounded uniformly continuous functions $v : X \to \mathbb{C}$ such that $\lim_{r\to\infty} v(ra+x)$ exists, for each $a \in X$, locally uniformly in $x \in X$. This algebra is larger than $\mathcal{E}(X)$. It would be interesting to find an explicit description of its spectrum.

We may now state our main results: they are consequences of the general theory developed so far (Section 1, Proposition 1, Theorem 2, and Example 2).

Theorem 3. *Let H be a self-adjoint operator on $L^2(X)$ affiliated to $\mathcal{E}(X) \rtimes X$. Then* s-$\lim_{r\to+\infty} T_{ra}HT_{ra}^* =: \tau_{\widehat{a}}(H)$ *exists $\forall a \in X \setminus \{0\}$ and $\sigma_{\rm ess}(H) = \overline{\cup}_{\alpha\in\mathbb{S}_X}\sigma(\tau_\alpha(H))$.*

Theorem 4. *Let h be as in Example 2 and $V = \sum V_Y$ with $V_Y : \mathcal{H}^s \to \mathcal{H}^{-s}$ symmetric operators such that $V_Y = 0$ but for a finite number of Y and satisfying:*

(1) $\exists\, \mu_Y, \nu_Y \geq 0$ with $\sum_Y \mu_Y < 1$ such that $\pm V_Y \leq \mu_Y h(P) + \nu_Y$,
(2) $\lim_{p\to 0} \|[S_p, V_Y]\|_{\mathcal{H}^s\to\mathcal{H}^{-s}} = 0$,
(3) $[T_y, V_Y] = 0$ for all $y \in Y$,
(4) $\tau_\alpha(V_Y) :=$ s-$\lim_{a\to\alpha} T_a V_Y T_a^*$ *exists in $B(\mathcal{H}^s,\mathcal{H}^{-s})$ for all $\alpha \in \mathbb{S}_{X/Y}$.*

Then the maps $\mathcal{H}^s \to \mathcal{H}^{-s}$ given by $h(P)+V$ and $h(P)+\sum_Y \tau_\alpha(V_Y)$ induce self-adjoint operators H and $\tau_\alpha(H)$ in $L^2(X)$ affiliated to $\mathscr{E}(X)$ and $\sigma_{\rm ess}(H) = \overline{\cup}_{\alpha\in\mathbb{S}_X}\sigma(\tau_\alpha(H))$.

Using also some techniques from [2], we obtain:

Example 3. Theorem 4 covers operators $H = \sum_{|k|,|\ell|\leq s} P^k a_{k\ell} P^\ell$ which are uniformly elliptic with the $a_{k\ell}$ finite sums of functions of the form $v_Y \circ \pi_Y$ with $v_Y : X/Y \to \mathbb{R}$ bounded measurable and such that $\lim_{z\to\alpha} v_Y(z)$ exists uniformly in $\alpha \in \mathbb{S}_{X/Y}$. The fact that we allow $a_{k\ell}$ to be only bounded measurable in the principal part of the operator (i. e. the terms with $|k| = |\ell| = s$) is not trivial.

In addition to the above mentioned results, we also use general results on cross-product C^*-algebras, their ideals, and their representations [4, 5, 13]. The maximal ideal spectrum of the algebra $\mathcal{E}(X)$ is of independent interest and can be used to study the regularity properties of the eigenvalues of the N-body Hamiltonian [1]. Its relation to the constructions of Vasy in [14] will be studied elsewhere.

Acknowledgements. We thank Bernd Ammann for several useful discussions.

REFERENCES

[1] B. Ammann, C. Carvalho, and V. Nistor. Regularity for eigenfunctions of Schrödinger operators. *Lett. Math. Phys.*, 101:49–84, 2012.
[2] M. Damak, and V. Georgescu. Self-adjoint operators affiliated to C^*-algebras, *Rev. Math. Phys.*, 16(2):257–280, 2004.
[3] J. Dereziński and C. Gérard. *Scattering theory of classical and quantum N-particle systems*. Texts and Monographs in Physics. Springer-Verlag, Berlin, 1997.
[4] J. Dixmier. *C^*-algebras*. North-Holland Publishing Co., Amsterdam, 1977.
[5] T. Fack and G. Skandalis. Sur les représentations et idéaux de la C^*-algèbre d'un feuilletage. *J. Operator Theory*, 8(1):95–129, 1982.
[6] V. Georgescu and A. Iftimovici. Crossed products of C^*-algebras and spectral analysis of quantum Hamiltonians. *Comm. Math. Phys.*, 228(3):519–560, 2002.
[7] V. Georgescu, and A. Iftimovici. Localizations at infinity and essential spectrum of quantum Hamiltonians. I. General theory. *Rev. Math. Phys.*, 18(4):417–483, 2006.
[8] B. Helffer and A. Mohamed. Caractérisation du spectre essentiel de l'opérateur de Schrödinger avec un champ magnétique. *Ann. Inst. Fourier (Grenoble)*, 38(2):95–112, 1988.

[9] Y. Last, Y. and B. Simon. The essential spectrum of Schrödinger, Jacobi, and CMV operators. *J. Anal. Math.*, 98183–220, 2006.
[10] R. Lauter, B. Monthubert, and V. Nistor. Pseudodifferential analysis on continuous family groupoids. *Doc. Math.*, 5:625–655 (electronic), 2000.
[11] A. Mageira. Graded C^*-algebras. *J. Funct. Anal.*, 2546:1683–1701, 2008.
[12] V. Rabinovich, S. Roch, and B. Silbermann. *Limit operators and their applications in operator theory*. volume 150 of *Operator Theory: Advances and Applications*. Birkhäuser Verlag, Basel, 2004.
[13] J. Renault. *A groupoid approach to C^*-algebras*, volume 793 of Lecture Notes in Mathematics. Springer, Berlin, 1980.
[14] A. Vasy. Propagation of singularities in many-body scattering. *Ann. Sci. École Norm. Sup. (4)*, 34(3):313–402, 2001.

V. GEORGESCU, DÉPARTEMENT DE MATHÉMATIQUES, UNIVERSITÉ DE CERGY-PONTOISE, 95000 CERGY-PONTOISE, FRANCE

E-mail address: vladimir.georgescu@math.cnrs.fr

V. NISTOR, UNIVERSITÉ DE LORRAINE, UFR MIM, ILE DU SAULCY, CS 50128, 57045 METZ CEDEX 01, AND, PENNSYLVANIA STATE UNIVERSITY, MATH. DEPT., UNIVERSITY PARK, PA 16802, USA

E-mail address: victor.nistor@univ-lorraine.fr